

# University of British Columbia CPSC 314 Computer Graphics Jan-Apr 2016

Tamara Munzner

## **Transformations 2**


http://www.ugrad.cs.ubc.ca/~cs314/Vjan2016

## Readings for Transformations 1-5

- Shirley/Marschner
  - Ch 6: Transformation Matrices
    - except 6.1.6, 6.3.1
  - Sect 12.2 Scene Graphs
- Gortler
  - Ch 2: Linear, Sec 2.5-2.6
  - Ch 3: Affine
  - Ch 4: Respect
  - Ch 5: Frames in Graphics, 5.3-5.4

## **2D Transformations**

### **2D Translation**



#### matrix multiplication

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

scaling matrix

#### vector addition

$$\begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} x + a \\ y + b \end{bmatrix} = \begin{bmatrix} x' \\ y' \end{bmatrix}$$

#### matrix multiplication

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

rotation matrix

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x' \\ y' \end{bmatrix}$$

## **Linear Transformations**

- linear transformations are combinations of
  - shear
  - scale
  - rotate
  - reflect

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

$$x' = ax + by$$

$$x' = ax + by$$
$$y' = cx + dy$$

- properties of linear transformations
  - satisfies T(sx+ty) = s T(x) + t T(y)
  - origin maps to origin
  - lines map to lines
  - parallel lines remain parallel
  - ratios are preserved
  - closed under composition

# Challenge

- matrix multiplication
  - for everything except translation
  - can we just do everything with multiplication?
    - then could just do composition, no special cases

# **Homogeneous Coordinates**

- represent 2D coordinates (x,y) with 3-vector (x,y,1)
  - use 3x3 matrices for 2D transformations


$$\mathbf{R}otation = \begin{bmatrix} \cos(\theta) & -\sin(\theta) & 0 \\ \sin(\theta) & \cos(\theta) & 0 \\ 0 & 0 & 1 \end{bmatrix} \qquad \mathbf{S}cale = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\mathbf{T}rans \, lation = \begin{bmatrix} 1 & 0 & T_x \\ 0 & 1 & T_y \\ 0 & 0 & 1 \end{bmatrix} \quad \text{use rightmost column}$$

$$\begin{bmatrix} 1 & 0 & a \\ 0 & 1 & b \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} x*1+a*1 \\ y*1+b*1 \\ 1 \end{bmatrix} = \begin{bmatrix} x+a \\ y+b \\ 1 \end{bmatrix}$$

## **Homogeneous Coordinates Geometrically**

point in 2D cartesian




## **Homogeneous Coordinates Geometrically**

#### homogeneous

#### cartesian

$$(x, y, w) \xrightarrow{/w} (\frac{x}{w}, \frac{y}{w})$$




- point in 2D cartesian + weight w = point P in 3D homog. coords
- multiples of (x,y,w)
  - form a line L in 3D
  - all homogeneous points on L represent same 2D cartesian point
  - example: (2,2,1) = (4,4,2) = (1,1,0.5)

## **Homogeneous Coordinates Geometrically**

#### homogeneous

#### cartesian

$$(x, y, w) \xrightarrow{/w} (\frac{x}{w}, \frac{y}{w})$$



- homogenize to convert homog. 3D point to cartesian 2D point:
  - divide by w to get (x/w, y/w, 1)
  - projects line to point onto w=1 plane
  - like normalizing, one dimension up
  - when w=0, consider it as direction
    - points at infinity
    - these points cannot be homogenized
    - lies on x-y plane
  - (0,0,0) is undefined

### **Affine Transformations**

- affine transforms are combinations of
  - linear transformations
  - translations

$$\begin{bmatrix} x' \\ y' \\ w \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$

- properties of affine transformations
  - origin does not necessarily map to origin
  - lines map to lines
  - parallel lines remain parallel
  - ratios are preserved
  - closed under composition

# **Homogeneous Coordinates Summary**

- may seem unintuitive, but they make graphics operations much easier
- allow all affine transformations to be expressed through matrix multiplication
  - we'll see even more later...
- use 3x3 matrices for 2D transformations
  - use 4x4 matrices for 3D transformations