



The Rendering Pipeline – Transformations

Wolfgang Heidrich

Wolfgang Heidrich



Course News

Assignment 0

- Due today!

Assignment 1

- Will be out Wednesday
- Due January 31

Homework 1

- Exercise problems for transformations
- Discussed in labs next week


Reading

- Chapter 5

Wolfgang Heidrich



The Rendering Pipeline



Modeling and Viewing Transformation

Affine transformations

- Linear transformations + translations
- Can be expressed as a 3x3 matrix + 3 vector

$$\mathbf{x}' = \mathbf{M} \cdot \mathbf{x} + \mathbf{t}$$


Scaling

Scaling

- A coordinate means multiplying each of its components by a scalar

Uniform scaling


- This scalar is the same for all components:



Scaling

Non-uniform scaling:

- different scalars per component:



How can we represent this in matrix form?



Scaling (2D)

Scaling operation:
$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} ax \\ by \end{pmatrix}$$

Or, in matrix form:
$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \underbrace{\begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}}_{\text{scaling matrix}} \cdot \begin{pmatrix} x \\ y \end{pmatrix}$$

Wolfgang Heidrich



Scaling (3D)


Scaling operation:
$$\begin{pmatrix} x' \\ y' \\ z' \end{pmatrix} = \begin{pmatrix} ax \\ by \\ cz \end{pmatrix}$$

Or, in matrix form:
$$\begin{pmatrix} x' \\ y' \\ z' \end{pmatrix} = \begin{pmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{pmatrix} \cdot \begin{pmatrix} x \\ y \\ z \end{pmatrix}$$

Wolfgang Heidrich



2D Rotation From Trig Identities



$$\begin{aligned}x &= r \cos(\phi) \\y &= r \sin(\phi) \\x' &= r \cos(\phi + \theta) \\y' &= r \sin(\phi + \theta)\end{aligned}$$

Trig Identity...

$$\begin{aligned}x' &= r \cos(\phi) \cos(\theta) - r \sin(\phi) \sin(\theta) \\y' &= r \sin(\phi) \cos(\theta) + r \cos(\phi) \sin(\theta)\end{aligned}$$

Substitute...

$$\begin{aligned}x' &= x \cos(\theta) - y \sin(\theta) \\y' &= x \sin(\theta) + y \cos(\theta)\end{aligned}$$

Wolfgang Heidrich



2D Rotation Matrix

Easy to capture in matrix form:

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{pmatrix} \cdot \begin{pmatrix} x \\ y \end{pmatrix}$$

Even though $\sin(q)$ and $\cos(q)$ are nonlinear functions of q ,

- x' is a linear combination of x and y
- y' is a linear combination of x and y

Wolfgang Heidrich



3D Rotation

- About x axis:

$$\begin{pmatrix} x' \\ y' \\ z' \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos(\theta) & -\sin(\theta) \\ 0 & \sin(\theta) & \cos(\theta) \end{pmatrix} \cdot \begin{pmatrix} x \\ y \\ z \end{pmatrix}$$

- About y axis:

$$\begin{pmatrix} x' \\ y' \\ z' \end{pmatrix} = \begin{pmatrix} \cos(\theta) & 0 & \sin(\theta) \\ 0 & 1 & 0 \\ -\sin(\theta) & 0 & \cos(\theta) \end{pmatrix} \cdot \begin{pmatrix} x \\ y \\ z \end{pmatrix}$$

- About z axis:

$$\begin{pmatrix} x' \\ y' \\ z' \end{pmatrix} = \begin{pmatrix} \cos(\theta) & -\sin(\theta) & 0 \\ \sin(\theta) & \cos(\theta) & 0 \\ 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} x \\ y \\ z \end{pmatrix}$$

Wolfgang Heidrich




Shear

Shear along x axis

- push points to right in proportion to height

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} ? & ? \\ ? & ? \end{pmatrix} \cdot \begin{pmatrix} x \\ y \end{pmatrix}$$



Wolfgang Heidrich




Shear

Shear along x axis

- push points to right in proportion to height

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} 1 & sh_x \\ 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} x \\ y \end{pmatrix}$$



Wolfgang Heidrich




Reflection

Reflect across x axis

- Mirror

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} ? & ? \\ ? & ? \end{pmatrix} \cdot \begin{pmatrix} x \\ y \end{pmatrix}$$



Wolfgang Heidrich




Reflection

Reflect across x axis

- Mirror

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \cdot \begin{pmatrix} x \\ y \end{pmatrix}$$



Wolfgang Heidrich




Affine Transformations

Translation:

- Add a constant (2D or 3D) vector:

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} t_x \\ t_y \end{pmatrix}$$



Wolfgang Heidrich

Compositing of Affine Transformations



In general:

- Transformation of geometry into coordinate system where operation becomes simpler
- Perform operation
- Transform geometry back to original coordinate system

Wolfgang Heidrich

Compositing of Affine Transformations



Example: 2D rotation around arbitrary center

- Consider this transformation

$$\mathbf{x}' = \underbrace{\mathbf{Id} \cdot \left(\overbrace{R(\phi)}^{\text{rotate by } \phi} \cdot \underbrace{(\mathbf{Id} \cdot \mathbf{x} - \mathbf{t})}_{\text{translate by } -\mathbf{t}} \right)}_{\text{translate by } \mathbf{t}} + \mathbf{t}$$

- i.e:

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} 1 & \\ & 1 \end{pmatrix} \cdot \left(\begin{pmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{pmatrix} \cdot \left(\begin{pmatrix} 1 & \\ & 1 \end{pmatrix} \cdot \begin{pmatrix} x \\ y \end{pmatrix} - \begin{pmatrix} a \\ b \end{pmatrix} \right) \right) + \begin{pmatrix} a \\ b \end{pmatrix}$$

Wolfgang Heidrich

Compositing of Affine Transformations



Composite transformation:

- Note that this is again an affine transformation

$$\begin{aligned}\mathbf{x}' &= \mathbf{Id} \cdot (R(\phi) \cdot (\mathbf{Id} \cdot \mathbf{x} - \mathbf{t})) + \mathbf{t} \\ &= \mathbf{Id} \cdot (R(\phi) \cdot \mathbf{x} - R(\phi) \cdot \mathbf{t}) + \mathbf{t} \\ &= R(\phi) \cdot \mathbf{x} + (R(\phi) \cdot (-\mathbf{t}) + \mathbf{t}) \\ &= R(\phi) \cdot \mathbf{x} + \mathbf{t}'\end{aligned}$$

This holds in general!

- All composites of affine transformations are themselves affine transformations!

Wolfgang Heidrich

Compositing of Affine Transformations



Two different interpretations of composite:


- 1) read from inside-out as transformation of object
 - 1a) Translate object by $-\mathbf{t}$
 - 1b) Rotate object by Φ
 - 1c) Translate object by \mathbf{t}
- 2) read from outside-in as transformation of the coordinate frame
 - 2c) Translate frame by \mathbf{t}
 - 2b) Rotate frame by $-\Phi$ (i.e. rotate object by Φ)
 - 2a) Translate frame by $-\mathbf{t}$

Wolfgang Heidrich

Compositing of Affine Transformations



Example scene:




Wolfgang Heidrich

Compositing of Affine Transformations



First Interpretation:

- Step 1: translate object by $-t$ (move to origin)




Wolfgang Heidrich

Compositing of Affine Transformations



First Interpretation:

- Step 2: rotate object by Φ




Wolfgang Heidrich

Compositing of Affine Transformations



First Interpretation:

- Step 3: translate object by t (move back)




Our composite example is a rotation around an arbitrary 2D point with position t !

Wolfgang Heidrich

Compositing of Affine Transformations



Example scene, second interpretation:




Wolfgang Heidrich

Compositing of Affine Transformations



Second interpretation:

- Step 1: translate frame (move origin to object)




Wolfgang Heidrich

Compositing of Affine Transformations



Second interpretation:

- Step 2: rotate frame by $-\Phi$ (i.e. rotate obj. by Φ)




Wolfgang Heidrich

Compositing of Affine Transformations



Second interpretation:

- Step 3: translate frame back (vector $-t$ in new frame!)



Wolfgang Heidrich

Compositing of Affine Transformations



NOTES:

- All transformations are **always with respect to the current coordinate frame**
- The results of both interpretations are **identical**
 - *Note that the object has the same relative position and orientation with respect to the coordinate frame!*

Wolfgang Heidrich

Compositing of Affine Transformations



Another Example: 3D rotation around arbitrary axis

- Rotate axis to z-axis
- Rotate by ϕ around z-axis
- Rotate z-axis back to original axis
- Composite transformation:

$$\begin{aligned} R(v, \phi) &= R_z^{-1}(\alpha) \cdot R_y^{-1}(\beta) \cdot R_z(\phi) \cdot R_y(\beta) \cdot R_z(\alpha) \\ &= (R_y(\beta) \cdot R_z(\alpha))^{-1} \cdot R_z(\phi) \cdot (R_y(\beta) \cdot R_z(\alpha)) \end{aligned}$$

Wolfgang Heidrich

Properties of Affine Transformations



Definition:

- A linear combination of points or vectors is given as

$$\mathbf{x} = \sum_{i=1}^n a_i \cdot \mathbf{x}_i, \text{ for } a_i \in \mathfrak{R}$$

- An affine combination of points or vectors is given as

$$\mathbf{x} = \sum_{i=1}^n a_i \cdot \mathbf{x}_i, \text{ with } \sum_{i=1}^n a_i = 1$$

Wolfgang Heidrich


Properties of Affine Transformations



Example:

- Affine combination of 2 points

$$\begin{aligned} \mathbf{x} &= a_1 \cdot \mathbf{x}_1 + a_2 \cdot \mathbf{x}_2, \text{ with } a_1 + a_2 = 1 \\ &= (1 - a_2) \cdot \mathbf{x}_1 + a_2 \cdot \mathbf{x}_2 \\ &= \mathbf{x}_1 + a_2 \cdot (\mathbf{x}_2 - \mathbf{x}_1) \end{aligned}$$




Wolfgang Heidrich

Properties of Affine Transformations



Definition:

- A convex combination is an affine combination where all the weights a_i are positive
- Note: this implies $0 \leq a_i \leq 1, i=1 \dots n$



Wolfgang Heidrich

Properties of Affine Transformations




Example:

- Convex combination of 3 points

$$\mathbf{x} = \alpha \cdot \mathbf{x}_1 + \beta \cdot \mathbf{x}_2 + \gamma \cdot \mathbf{x}_3$$

$$\text{with } \alpha + \beta + \gamma = 1, 0 \leq \alpha, \beta, \gamma \leq 1$$

- $\alpha, \beta,$ and γ are called *Barycentric coordinates*



Wolfgang Heidrich

Properties of Affine Transformations



Theorem:

- The following statements are synonymous
 - A transformation $T(x)$ is affine, i.e.:

$$\mathbf{x}' = T(\mathbf{x}) := \mathbf{M} \cdot \mathbf{x} + \mathbf{t},$$

for some matrix \mathbf{M} and vector \mathbf{t}

- $T(x)$ preserves affine combinations, i.e.

$$T\left(\sum_{i=1}^n a_i \cdot \mathbf{x}_i\right) = \sum_{i=1}^n a_i \cdot T(\mathbf{x}_i), \text{ for } \sum_{i=1}^n a_i = 1$$

- $T(x)$ maps parallel lines to parallel lines


Wolfgang Heidrich

Properties of Affine Transformations



Preservation of affine combinations:

- Can compute transformation of every point on line or triangle by simply transforming the *control points*



Wolfgang Heidrich



Coming Up

This week:

- Affine Transformations with Homogeneous Coordinates
- Transformation Hierarchies

Next week

- Perspective Transformations