



University of
British Columbia



Chapter 4

Transformations






Transformations

- Transformation = *one-to-one* and *onto* mapping of R^n to itself
- *Affine* transformation – $T(v) = Av+b$
 - A – matrix
 - v, b – vectors


$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}$$






Geometric Transformations

- *Geometric Transformation* = affine transformation with geometric meaning



- Mathematically transformations are defined on vectors \Rightarrow for point P, use vector P-Origin





Matrix Representation

- Combining transformations


$$ST(v) = C(Av + b) + d = (CA)v + (Cb + d)$$

- Same format (multiply by matrix & add vector)

$$A' = CA \quad b' = Cb + d$$

- Can reduce to matrix multiplication only
 - Homogeneous coordinates (later on)






Transformations – sub topics




- 2D Transformations
- Homogeneous Coordinates
- 3D Transformations
- Composing Transformations
- Transformation Hierarchies
- Transforming Normals
- Assignment 1 – dinosaurs
 - Use transformations to create and animate dinosaurs made from spheres






Transformations

- Transforming an object = transforming all its points
- Transforming a polygon = transforming its vertices
 - Why?




Applications

- Viewing (more later)





link	mass (kg)
L1	0.15
L2	0.10
L3	0.30



- Modeling

- Articulation








Scaling

- $V = \begin{pmatrix} v_x \\ v_y \end{pmatrix}$ – vector in XY plane
- *Scaling* operator S with parameters (s_x, s_y) :

$$S^{(s_x, s_y)}(V) = \begin{pmatrix} s_x v_x \\ s_y v_y \end{pmatrix}$$





Scaling



- Matrix form:

$$S^{(s_x, s_y)}(V) = \begin{pmatrix} s_x & 0 \\ 0 & s_y \end{pmatrix} \begin{pmatrix} v_x \\ v_y \end{pmatrix} = \begin{pmatrix} s_x v_x \\ s_y v_y \end{pmatrix}$$

- Independent in x and y



Rotation




- Polar form:

$$v = \begin{pmatrix} v_x \\ v_y \end{pmatrix} = \begin{pmatrix} r \cos \alpha \\ r \sin \alpha \end{pmatrix}$$

- Rotating v counterclockwise by θ to w:

$$w = \begin{pmatrix} w_x \\ w_y \end{pmatrix} = \begin{pmatrix} r \cos(\alpha + \theta) \\ r \sin(\alpha + \theta) \end{pmatrix} = \begin{pmatrix} r \cos \alpha \cos \theta - r \sin \alpha \sin \theta \\ r \cos \alpha \sin \theta + r \sin \alpha \cos \theta \end{pmatrix}$$





Rotation


- Matrix form:

$$w = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} r \cos \alpha \\ r \sin \alpha \end{pmatrix} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} v$$

- *Rotation* operator R (at the origin) with parameter θ :

$$R^\theta = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$





Rotation Properties


- R^θ is orthonormal

$$(R^\theta)^{-1} = (R^\theta)^T$$


- $R^{-\theta}$ - rotation by $-\theta$ is

$$R^{-\theta} = \begin{bmatrix} \cos(-\theta) & -\sin(-\theta) \\ \sin(-\theta) & \cos(-\theta) \end{bmatrix} = \begin{bmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{bmatrix} = (R^\theta)^{-1}$$






Translation



- Translation operator T with parameters (t_x, t_y) :

$$T^{(t_x, t_y)}(v) = \begin{pmatrix} v_x + t_x \\ v_y + t_y \end{pmatrix}$$






Translation - Homogeneous Coordinates

- To represent T in matrix form – introduce homogeneous coordinates:

$$v = \begin{pmatrix} v_x \\ v_y \end{pmatrix} \rightarrow v^h = \begin{pmatrix} v_x^h \\ v_y^h \\ v_w^h \end{pmatrix} = \begin{pmatrix} v_x \\ v_y \\ 1 \end{pmatrix}$$





Translation - Homogeneous Coordinates


- Conversion (projection) from homogeneous space to Euclidean:

$$v = \begin{pmatrix} v_x \\ v_y \end{pmatrix} = \begin{pmatrix} v_x^h / v_w^h \\ v_y^h / v_w^h \end{pmatrix}$$

- Projections is not 1:1

$$\begin{pmatrix} 2 \\ 2 \\ 1 \end{pmatrix}, \begin{pmatrix} 4 \\ 4 \\ 2 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 0.5 \end{pmatrix} \text{ all project to } \begin{pmatrix} 2 \\ 2 \end{pmatrix}$$






Translation

- Using homogeneous coordinates, translation operator may be expressed as:

$$T^{(t_x, t_y)}(v^h) = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{pmatrix} v_x \\ v_y \\ 1 \end{pmatrix} = \begin{pmatrix} v_x + t_x \\ v_y + t_y \\ 1 \end{pmatrix}$$






Homogeneous Coordinates

$$\text{Rotation} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) & 0 \\ \sin(\theta) & \cos(\theta) & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\text{Scale} = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & 1 \end{bmatrix}$$


Other ideas for uniform scale?






Transformation Composition

- What operation rotates XY by θ around $P = \begin{pmatrix} p_x \\ p_y \end{pmatrix}$?
- Answer:
 - Translate P to origin
 - Rotate around origin by θ
 - Translate back






Transformation Composition


$$T^{(p_x, p_y)} R^\theta T^{(-p_x, -p_y)}(V)$$

$$= \begin{bmatrix} 1 & 0 & p_x \\ 0 & 1 & p_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & -p_x \\ 0 & 1 & -p_y \\ 0 & 0 & 1 \end{bmatrix} \begin{pmatrix} v_x \\ v_y \\ 1 \end{pmatrix}$$





Transformations Quiz



- What do these transformations do?


$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$$


$$\begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & a \\ 0 & 1 \end{bmatrix}$$





Transformations Quiz




- And these homogeneous ones?


$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0.5 \end{bmatrix}$$







Transformations Quiz



- How to mirror through arbitrary line in XY?
- What transformation achieves this?





Shear & Mirroring/Reflection


- Shear (canonical)


$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & a \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ a & 1 \end{bmatrix}$$

- Mirroring/Reflection

$$\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$$

- What is the relation between shears and rotations?





Linear Transformations


- Combinations of
 - shear
 - scale
 - rotate
 - reflect
- Properties (why?)
 - satisfies $T(sx+ty) = s T(x) + t T(y)$
 - origin maps to origin
 - Straight lines map to straight lines
 - parallel lines remain parallel
 - closed under composition

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

$$x' = ax + by$$

$$y' = cx + dy$$






Affine Transformations

- Combinations of
 - linear transformations
 - translations
- Properties (why?)
 - origin does not necessarily map to origin
 - lines map to lines
 - parallel lines remain parallel
 - ratios are preserved
 - closed under composition

$$\begin{bmatrix} x' \\ y' \\ w \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$





3D Transformations

- All 2D transformations extend to 3D
- In homogeneous coordinates:

Scaling

$$S^{(s_x, s_y, s_z)} = \begin{bmatrix} s_x & 0 & 0 & 0 \\ 0 & s_y & 0 & 0 \\ 0 & 0 & s_z & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Translation

$$T^{(t_x, t_y, t_z)} = \begin{bmatrix} 1 & 0 & 0 & t_x \\ 0 & 1 & 0 & t_y \\ 0 & 0 & 1 & t_z \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Rotation around the z axis

$$R_z^\theta = \begin{bmatrix} \cos \theta & -\sin \theta & 0 & 0 \\ \sin \theta & \cos \theta & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

glScalef(a,b,c);


glTranslatef(a,b,c);

glRotatef(angle,0,0,1);



University of
British Columbia





3D Rotation in X, Y

around x axis:

$$\begin{bmatrix} x' \\ y' \\ z' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\theta & -\sin\theta & 0 \\ 0 & \sin\theta & \cos\theta & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix}$$

around y axis:

$$\begin{bmatrix} x' \\ y' \\ z' \\ 1 \end{bmatrix} = \begin{bmatrix} \cos\theta & 0 & \sin\theta & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\theta & 0 & \cos\theta & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix}$$


glRotatef(angle,1,0,0);

glRotatef(angle,0,1,0);


- general OpenGL command

glRotatef(angle,x,y,z);






Arbitrary Rotation



- Problem:
 - Given two orthonormal coordinate systems XYZ and UVW
 - Find transformation from one to the other
- Answer:
 - Transformation matrix R whose columns are U, V, W :



$$R = \begin{bmatrix} u_x & v_x & w_x \\ u_y & v_y & w_y \\ u_z & v_z & w_z \end{bmatrix}$$




Arbitrary Rotation

- Proof:

$$R(X) = \begin{bmatrix} u_x & v_x & w_x \\ u_y & v_y & w_y \\ u_z & v_z & w_z \end{bmatrix} \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} u_x \\ u_y \\ u_z \end{pmatrix} = U$$

- Similarly $R(Y) = V$ & $R(Z) = W$






Arbitrary Rotation (cont.)

- Inverse (=transpose) transformation R^{-1} provides mapping from UVW to XYZ
- E.g.

$$R^{-1}(U) = \begin{bmatrix} u_x & u_y & u_z \\ v_x & v_y & v_z \\ w_x & w_y & w_z \end{bmatrix} \begin{pmatrix} u_x \\ u_y \\ u_z \end{pmatrix} = \begin{pmatrix} u_x^2 + u_y^2 + u_z^2 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = X$$

- Comment: Used for mapping between XY and arbitrary plane





3D Shear

- shear in x

$$x\text{shear}(sy, sz) = \begin{bmatrix} 1 & sy & sz & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$


- shear in y

$$y\text{shear}(sx, sz) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ sx & 1 & sz & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

- shear in z

$$z\text{shear}(sx, sy) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ sx & sy & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$





Undoing Transformations: Inverses

$$\mathbf{T}(x, y, z)^{-1} = \mathbf{T}(-x, -y, -z)$$

$$\mathbf{T}(x, y, z) \mathbf{T}(-x, -y, -z) = \mathbf{I}$$


$$\mathbf{R}(z, \theta)^{-1} = \mathbf{R}(z, -\theta) = \mathbf{R}^T(z, \theta) \quad (\mathbf{R} \text{ is orthogonal})$$

$$\mathbf{R}(z, \theta) \mathbf{R}(z, -\theta) = \mathbf{I}$$

$$\mathbf{S}(sx, sy, sz)^{-1} = \mathbf{S}\left(\frac{1}{sx}, \frac{1}{sy}, \frac{1}{sz}\right)$$

$$\mathbf{S}(sx, sy, sz) \mathbf{S}\left(\frac{1}{sx}, \frac{1}{sy}, \frac{1}{sz}\right) = \mathbf{I}$$





3D Transformations - Composition

- Questions:

- Is $S_1S_2 = S_2S_1$?

- Is $T_1T_2 = T_2T_1$?


- Is $R_1R_2 = R_2R_1$?

- Is $S_1R_2 = R_2S_1$?

-



University of
British Columbia



Composing Translations


$$T1 = T(dx_1, dy_1) = \begin{bmatrix} 1 & & dx_1 \\ & 1 & dy_1 \\ & & 1 \\ & & & 1 \end{bmatrix} \quad T2 = T(dx_2, dy_2) = \begin{bmatrix} 1 & & dx_2 \\ & 1 & dy_2 \\ & & 1 \\ & & & 1 \end{bmatrix}$$

$P'' = T2 \bullet P' = T2 \bullet [T1 \bullet P] = [T2 \bullet T1] \bullet P$, where

$$T2 \bullet T1 = \begin{bmatrix} 1 & & dx_1 + dx_2 \\ & 1 & dy_1 + dy_2 \\ & & 1 \\ & & & 1 \end{bmatrix}$$

Translations add





Composing Transformations

- scaling

$$S2 \bullet S1 = \begin{bmatrix} sx_1 * dx_2 & & \\ & sy_1 * sy_2 & \\ & & 1 \\ & & & 1 \end{bmatrix}$$

scales multiply

- rotation


$$R2 \bullet R1 = \begin{bmatrix} \cos(\theta_1 + \theta_2) & -\sin(\theta_1 + \theta_2) & & \\ \sin(\theta_1 + \theta_2) & \cos(\theta_1 + \theta_2) & & \\ & & 1 & \\ & & & 1 \end{bmatrix}$$

rotations add



Composing Transformations


ORDER MATTERS!



$T_a T_b = T_b T_a$, but $R_a R_b \neq R_b R_a$ and $T_a R_b \neq R_b T_a$



University of
British Columbia



Another Transformations Quiz



- What does each transformation preserve?

	lines	parallel lines	distance	angles	normals	convexity
scaling						
rotation						
translation						
shear						

