




Chapter 6

Geometric Modeling Part I: Terminology & Splines

An Example




Operations

Acquisition/Generation

Transmission/Storage



Editing



University of British Columbia

Geometry





- Mathematical models of real world objects shape
- Categories:
 - Boundary representations
 - Freeform
 - Meshes & subdivision
 - Volumetric representations
 - Primitive based
 - Voxels



University of British Columbia

Volumetric - Voxelization






- Voxel based
 - Space subdivided into equal size boxes – each has in/out flag
 - Common in imaging applications (CT,MRI,Ultrasound)
- Extension – Octree (recursive construction)
- To draw use *iso-surfaces* – boundary between voxels with different flag



University of British Columbia

Volumetric - Primitives

- Use set of volumetric primitives
 - Box, sphere, cylinder, cone, etc...
- For complex objects use boolean operations
 - Union
 - Intersection
 - Subtraction



University of British Columbia


Freeform Representation

- Explicit form: $z=z(x,y)$ Explicit is a special case of implicit and parametric form
- Implicit form: $f(x,y,z)=0$
- Parametric form: $[x(u,v),y(u,v),z(u,v)]$
- Example – origin centered sphere of radius R :

Explicit:
 $z = +\sqrt{R^2 - x^2 - y^2} \cup z = -\sqrt{R^2 - x^2 - y^2}$

Implicit:
 $x^2 + y^2 + z^2 - R^2 = 0$


Parametric:
 $(x,y,z) = (R \cos \theta \cos \psi, R \sin \theta \cos \psi, R \sin \psi), \theta \in [0, 2\pi], \psi \in [-\frac{\pi}{2}, \frac{\pi}{2}]$




University of British Columbia

Splines – Free Form Curves


- Description = basis functions + coefficients
- Geometric meaning of coefficients (base)
 - Approximate/interpolate set of positions, derivatives, etc..



- Usually parametric




University of British Columbia




Polynomial Bases

- Monomial basis
 - Geometrically meaningless $\{1, t, t^2, t^3, \dots\}$
- Other bases
 - Bezier
 - Hermitte
 - Lagrange, B-Spline, ...
- Number of coefficients = polynomial rank
- Geometric meaning
 - coefficients - positions/derivatives, etc...
- Advantage
 - easy to analyze, derivatives remain polynomial




University of
British Columbia





Lagrange

- Interpolates **control points**
 - P_0, \dots, P_n
- Base function $P_i(t)$ per control-point P_i
 - $P_i(t) = 1$ at i/n
 - $P_i(t) = 0$ at $j/n, j \neq i$
- Each base function is a polynomial of degree n
$$P_i(t) = a_0^i + a_1^i t + a_2^i t^2 + \dots + a_n^i t^n$$
- Uniqueness
 - n equations in n unknowns (a_j^i)




University of
British Columbia






Analytic Continuity

- $C_1(t)$ & $C_2(t)$, $t \in [0, 1]$ - parametric curves
- Level of continuity at $C_1(1)$ and $C_2(0)$ is:
 - C^{-1} : $C_1(1) \neq C_2(0)$ (discontinuous)
 - C^0 : $C_1(1) = C_2(0)$ (positional continuity)
 - C^k , $k > 0$: continuous up to k -th derivative




$C_1^{(j)}(1) = C_2^{(j)}(0), 0 \leq j \leq k$

- Continuity of single curve defined similarly
 - for polynomial bases C^∞




University of British Columbia




Geometric Continuity

- Analytic continuity - too strong a requirement
- Geometric continuity – common curve is geometrically smooth (per given level k)
 - G^k , $k \leq 0$: Same as C^k
 - G^k , $k = 1$: $C'_1(1) = \alpha C'_2(0)$ $\alpha > 0$
 - G^k , $k \geq 0$: In arc-length reparameterization of $C_1(t)$ & $C_2(t)$, the two are C^k




University of British Columbia




Geometric Continuity

- E.g.
 - $C_1(t) = [\cos(t), \sin(t)] \quad t \in [-0.5\pi, 0]$
 - $C_2(t) = [\cos(t), \sin(t)] \quad t \in [0, 0.5\pi]$
 - $C_3(t) = [\cos(2t), \sin(2t)] \quad t \in [0, 0.25\pi]$




C_2, C_3
 C_1

- $C_1(t) \text{ \& } C_2(t)$ are C^k (& G^k) continuous
- $C_1(t) \text{ \& } C_3(t)$, are G^k continuous (not C^k)



University of
British Columbia




Hermite Cubic Basis

- Geometrically-oriented basis for cubic polynomials
 - 2 positions + 2 tangents
- Has to satisfy

$$h_{i,j}(t): i, j = 0, 1, \quad t \in [0, 1]$$

curve	$h(0)$	$h(1)$	$h'(0)$	$h'(1)$
$h_{00}(t)$	1	0	0	0
$h_{01}(t)$	0	1	0	0
$h_{10}(t)$	0	0	1	0
$h_{11}(t)$	0	0	0	1



University of
British Columbia



Hermite Cubic Basis (cont'd)

- Four cubics which satisfy the conditions

$$h_{00}(t) = t^2(2t-3)+1 \quad h_{01}(t) = -t^2(2t-3)$$

$$h_{10}(t) = t(t-1)^2 \quad h_{11}(t) = t^2(t-1)$$

- Obtain - solve 4 linear equations in 4 unknowns for each basis function


University of British Columbia

Hermite Cubic Basis (cont'd)

- Let $C(t)$ be a cubic polynomial defined as the linear combination:


$$C(t) = P_0 h_{00}(t) + P_1 h_{01}(t) + T_0 h_{10}(t) + T_1 h_{11}(t)$$

- What are $C(0)$, $C(1)$, $C'(0)$, $C'(1)$?




hermite

- $C(0)=P_0$, $C(1) = P_1$, $C'(0)=T_0$, $C'(1)=T_1$
- To generate a curve through P_0 & P_1 with slopes T_0 & T_1 use

$$C(t) = P_0 h_{00}(t) + P_1 h_{01}(t) + T_0 h_{10}(t) + T_1 h_{11}(t)$$



University of British Columbia




Natural Cubic Splines

- Standard spline input – set of points $\{P_i\}_{i=0}^n$
 - No derivatives
- Interpolate by n cubic segments:
 - Derive $\{T_i\}_{i=0}^n$ from continuity constraints
 - Solve 4n equations

Interpolation ($2n$ equations):
 $C_i(0) = P_{i-1} \quad C_i(1) = P_i \quad i = 1, \dots, n$
 C^1 continuity constraints ($n - 1$ equations):
 $C'_i(1) = C'_{i+1}(0) \quad i = 1, \dots, n - 1$
 C^2 continuity constraints ($n - 1$ equations):
 $C''_i(1) = C''_{i+1}(0) \quad i = 1, \dots, n - 1$





University of
British Columbia




Natural Cubic Splines

- Need another 2 equations to reach 4n
- Options
 - Natural end conditions: $C''_1(0) = 0, C''_n(1) = 0$
 - Prescribed end conditions
 (derivative available): $C'_1(0) = T_0, C'_n(1) = T_n$






University of
British Columbia




B-Splines


- Idea: Generate basis where functions are continuous cross domains



- Control point controls set of basis functions (to preserve continuity)
- Alternative view: continuous basis functions defined on several domains



University of
British Columbia





Uniform Cubic B-Spline Curves

- Definition


$$C(t) = \sum_{i=0}^{n-1} P_i N_i^3(t) \quad t \in [3, n]$$

$$N_i^3(t) = \begin{cases} r^3 / 6 & r = t - i \quad t \in [i, i+1] \\ (-3r^3 + 3r^2 + 3r + 1) / 6 & r = t - i - 1 \quad t \in [i+1, i+2] \\ (3r^3 - 6r^2 + 4) / 6 & r = t - i - 2 \quad t \in [i+2, i+3] \\ (1-r)^3 / 6 & r = t - i - 3 \quad t \in [i+3, i+4] \end{cases}$$







University of
British Columbia




Uniform Cubic B-Spline Curves

- For any $t \in [3, n]$ $\sum_{i=j-3}^j N_i^3(t) = 1$
- For any $t \in [j, j+1]$ only 4 basis functions are non zero
- Any point on cubic B-Spline is affine combination of at most 4 control points





University of British Columbia




Boundary Conditions for B-Splines

- B-Splines do not interpolate any control points
 - in particular end points
- Way to force endpoint interpolation:
 - Let $P_0 = P_1 = P_2$ and same for other end
- Question:
 - What is the shape of the curve at endpoints if this method is used ?





University of British Columbia




B-Spline Curve Properties

- For n control points, $C(t)$ is a piecewise polynomial of degree 3, defined for
- $C(t) \in \bigcup_{i=0}^{n-3} CH(P_i, \dots, P_{i+3}) \quad t \in [3, n]$
- $C(t)$ is affine invariant
- Questions:
 - What is $C(i)$?
 - What is $C'(i)$?
 - What is the continuity of $C(t)$?






bspline



University of
British Columbia



NURBs


- B-Spline

$$C(t) = \sum_{i=0}^{n-1} P_i N_i^3(t) \quad t \in [3, n]$$


$$N_i^3(t) = \begin{cases} r^3 / 6 & r = t - i & t \in [i, i + 1] \\ (-3r^3 + 3r^2 + 3r + 1) / 6 & r = t - i - 1 & t \in [i + 1, i + 2] \\ (3r^3 - 6r^2 + 4) / 6 & r = t - i - 2 & t \in [i + 2, i + 3] \\ (1 - r)^3 / 6 & r = t - i - 3 & t \in [i + 3, i + 4] \end{cases}$$

- Non-Uniform – different interval lengths (knots)
- Rational – rational basis functions

$$C(t) = \frac{\sum_{i=0}^{n-1} w_i P_i N_i^3(t)}{\sum_{i=0}^{n-1} w_i N_i^3(t)} \quad t \in [3, n]$$



University of
British Columbia




From Curves to Surfaces


- Curve is expressed as inner product of P_i coefficients and basis functions

$$C(u) = \sum_{i=0}^n P_i B_i(u)$$
- To extend curves to surfaces - treat surface as a curve of curves
- Assume P_i is not constant, but a function of second parameter v : $P_i(v) = \sum_{j=0}^m Q_{ij} B_j(v)$


$$C(u,v) = \sum_{i=0}^n \sum_{j=0}^m Q_{ij} B_j(v) B_i(u)$$



bezpatch



University of British Columbia




Bilinear Patches

- Bilinear interpolation of 4 3D points - 2D analog of 1D linear interpolation between 2 points in the plane
- Given $P_{00}, P_{01}, P_{10}, P_{11}$ associated parametric bilinear surface for $u, v \in [0,1]$ is:

$$P(u,v) = (1-u)(1-v)P_{00} + (1-u)vP_{01} + u(1-v)P_{10} + uvP_{11}$$

- Questions:
 - What does an isoparametric curve of a bilinear patch look like ?
 - When is a bilinear patch planar?



University of British Columbia